

Neutrino ground state in a dense star

Ken Kiers*

*High Energy Theory, Department of Physics
Brookhaven National Laboratory, Upton, NY 11973-5000, USA*

Michel H.G. Tytgat†

*High Energy Theory, Department of Physics
Brookhaven National Laboratory, Upton, NY 11973-5000, USA
and
Service de Physique Théorique
Université Libre de Bruxelles, CP225
Bd du Triomphe, 1050 Bruxelles, Belgium*

Abstract

It has recently been argued that long range forces due to the exchange of massless neutrinos give rise to a very large self-energy in a dense, finite-ranged, weakly charged medium. Such an effect, if real, would destabilize a neutron star. To address this issue we have studied the related problem of a massless neutrino field in the presence of an external, static electroweak potential of finite range. To be precise, we have computed to one loop the exact vacuum energy for the case of a spherical square well potential of depth α and radius R . For small wells, the vacuum energy is reliably determined by a perturbative expansion in the external potential. For large wells, however, the perturbative expansion breaks down. A manifestation of this breakdown is that the vacuum carries a non-zero neutrino charge. The energy and neutrino charge of the ground state are, to a good approximation for large wells, those of a neutrino condensate with chemical potential $\mu = \alpha$. Our results demonstrate explicitly that long-range forces due to the exchange of massless neutrinos do not threaten the stability of neutron stars.

*Email: kiers@bnl.gov

†Email: mtytgat@muon.phy.bnl.gov

I. INTRODUCTION

If neutrinos are massless, the exchange of neutrino-antineutrino pairs gives rise to long range interactions between weakly charged particles, such as neutrons [1,2]. On dimensional grounds, the potential between two neutrons separated by a distance r must be of the form

$$V \sim \frac{G_F^2}{r^5}$$

(where G_F is the Fermi constant), and thus falls off rapidly at large distances. As was first remarked by Feynman [3,4], an interesting effect may arise if there is a finite density of matter near the two neutrons. A 4-body process, for instance, can give rise to a potential that behaves like

$$V \sim \frac{G_F^4 N^2}{R^8} \frac{1}{r}, \quad (1.1)$$

if $R \gg r$, where N is the total number of particles in the cloud of matter and R is the distance between the matter and the two test neutrons. Because of the large N^2 enhancement factor in (1.1), the effective coupling $G_F^4 N^2 / R^8$ can be of order unity. In this case, far-away matter would give a large contribution to the two-body potential. Since this effect would be even more important for higher-order processes, one would then have to take into account diagrams of arbitrarily high order.

More recently, Fischbach [5] has studied the contribution to the energy of a neutron star due to the exchange of massless neutrinos. Large combinatoric enhancements, similar to those envisioned by Feynman, led him to conclude that a neutron star could not be gravitationally bound unless neutrinos are massive, $m_\nu \gtrsim 0.4$ eV.¹

Abada *et al* [9] have contended that Fischbach's calculation was incorrect. They have concluded that the neutrino contribution to the energy of a neutron star is essentially negligible. While the arguments given by Abada *et al* were compelling, the calculations upon which they were based did not take into account the finite size of the star. As we will demonstrate in this paper, the conclusions reached in [9] were nevertheless essentially correct.

We believe that we have completed the proof of the assertions made in [9]. We have studied the ground state properties of a massless neutrino field propagating in a finite-range electroweak potential, like the one induced by a finite density of neutrons. To be precise, we have considered a spherical square well electroweak potential of depth α and radius R , and have computed to one loop the exact energy and neutrino number of the ground state. By exact we mean that our results are non-perturbative in the external potential. In order to carry out this program, we have used an approach due initially to Schwinger [10].² For

¹ Smirnov and Vissani [6] have argued that the long range forces can also be screened by the neutrino condensate that resides in a star [7]. However, unlike screening at finite temperature, screening at finite density is not perfect. Fischbach, *et al* [8] have argued, in response, that in a finite system a residual effect could still blow up a star (or make it collapse into a black hole).

²Similar calculations are found in non-perturbative computations of quantum corrections to solitons [11–13] and in studies of QED in strong fields [14].

technical reasons, the potentials which we consider are of relatively modest strength and range compared to those of a realistic neutron star. As we shall argue, our calculation is nevertheless sufficient to demonstrate that the neutrino ground state energy in a neutron star does not vary wildly as the radius or density of the star is increased, but is in fact very well-behaved.

The paper is organized as follows. In Section II we introduce the reader to Fischbach's results and review the critique advanced by Abada, *et al.* We then make the case for our approach to the question. We show that we have a well-defined problem to which we can give a definitive solution.

Section III is essentially independent of the specific application to the neutron star problem and, we believe, is of interest in its own right. We first solve the Weyl equation in the presence of a spherical square well potential. Because the fermions are massless, there are no bound states but only scattering solutions. From the expression for the phase shifts, we compute the properly renormalized energy of the neutrino ground state in the presence of the potential. For small wells, $\alpha R \ll \pi$, we show that the energy of the ground state is perturbative in the potential. As the size of the potential increases, the perturbative expansion breaks down. A symptom of this breakdown is that for $\alpha R \geq \pi$ the ground state carries a non-zero neutrino charge. We show that, to a very good approximation for large wells, the ground state energy and neutrino number are those of a neutrino condensate with chemical potential $\mu = \alpha$, a result anticipated by Abada *et al.* That this result also holds for a neutron star follows trivially.

We emphasize that we have studied precisely the same problem as Fischbach did. Our approach, however, is quite different. We conclude that long range forces due to the exchange of massless neutrinos do not destabilize a star. There is indeed a shift in vacuum energy, but – maybe unfortunately – it is too tiny to be of any consequence regarding the fate of a neutron star.

Related lines of argument can be found in the recent literature. In [15], a 1+1 dimensional star is studied as a toy model. In this case, the vacuum energy can be calculated exactly and happens to vanish. It is also shown that the presence of a boundary implies the existence of a neutrino condensate. This point is also emphasized in [16]. Both papers stress that a complete 3 + 1 dimensional calculation for a system with a boundary is necessary in order to settle the issue definitively.

II. THE NEUTRINO GROUND STATE IN A NEUTRON STAR

A. Preliminaries

Before embarking on our main calculation, we introduce some useful concepts and formulas. The stated goal in Ref. [5] is to calculate the shift in vacuum energy due to neutrino exchange in a neutron star:

$$W = \langle \hat{0} | H | \hat{0} \rangle - \langle 0 | H_0 | 0 \rangle. \quad (2.1)$$

Here H denotes the Hamiltonian in the medium and $|\hat{0}\rangle$ refers to the true ground state of the system. The second term is the usual matter-free vacuum energy. For fermions, Eq. (2.1) amounts to

$$W = -\frac{1}{2} \left\{ \sum_{i>0} (E_i - E_i^0) - \sum_{i<0} (E_i - E_i^0) \right\}, \quad (2.2)$$

where the indices i refer to the positive and negative energy levels of the neutrinos. A first step toward the actual computation of (2.1) is the following expression due to Schwinger [10],

$$W = - \int d^3x \frac{\partial}{\partial t} \text{tr} \left[S_F(x, x') - S_F^0(x, x') \right]_{x' \rightarrow x} \quad (2.3)$$

where $S_F(x, x')$ is the 2×2 Feynman propagator for the massless left-handed neutrino field in the presence of the finite neutron density and $S_F^0(x, x')$ is that of the free field. In (2.3), the limit $x' \rightarrow x$ is taken symmetrically and averaged.

Let us now drop any explicit reference to the neutrons of the star and consider instead the following effective Lagrangian for the neutrino field

$$\mathcal{L}_{\text{eff}} = \bar{\psi}_L \left[i\not{\partial} + \alpha\gamma^0 \right] \psi_L \quad (2.4)$$

where $\psi_L = L\psi$, $L = (1 - \gamma_5)/2$, and where

$$\alpha(\vec{x}) = G_F \rho_n(\vec{x}) / \sqrt{2} \sim 20 \text{ eV}$$

is the electroweak potential induced by the finite neutron density ($\rho_n \approx 0.4 \text{ fm}^{-3}$ in a typical neutron star). The static potential $\alpha > 0$ is attractive for neutrinos and repulsive for antineutrinos. The effective Lagrangian of (2.4) has been derived in various ways from the underlying electroweak theory [17–19] and applied to studies of the Mikheyev-Smirnov-Wolfenstein (MSW) effect [20]. It provides us with a most convenient tool to study the coherent behavior of low-energy neutrinos in a neutron star, on scales large compared to the size of a neutron $r_n \sim 1 \text{ fm} \approx (200 \text{ MeV})^{-1}$.³

Since the potential is static, it is convenient to introduce

$$S_F^{(0)}(x, x') = \frac{1}{2\pi} \int_C d\omega e^{-i\omega(t-t')} G_{(0)}(\vec{x}, \vec{x}'; \omega) \quad (2.5)$$

with C representing the usual Feynman contour. The resolvents $G(\vec{x}, \vec{x}'; \omega)$ and $G_0(\vec{x}, \vec{x}'; \omega)$ satisfy

$$(\omega - H_{(0)}) G_{(0)}(\vec{x}, \vec{x}'; \omega) = -\delta^3(\vec{x} - \vec{x}') \quad (2.6)$$

with the Hamiltonian from (2.4)

$$H = -\alpha + i\vec{\sigma} \cdot \vec{\nabla}_x; \quad H_0 = i\vec{\sigma} \cdot \vec{\nabla}_x. \quad (2.7)$$

The Schwinger formula, (2.3), becomes

³ The fact that the effective theory, as in (2.4), is anomalous should not worry us. The underlying theory (the Standard Model) is perfectly well defined. Besides, for a static potential in $3 + 1$ dimensions, there are no anomalous effects because $\vec{E} \cdot \vec{B} = 0$.

$$\begin{aligned}
W &= \frac{i}{2\pi} \int_C d\omega \, \omega \, \text{Tr}_{\mathbf{x}} [G(\omega) - G_0(\omega)] \\
&= -\frac{i}{2\pi} \int_C d\omega \, \omega \, \text{Tr}_{\mathbf{x}} \left[\frac{1}{\omega - H} - \frac{1}{\omega - H_0} \right]
\end{aligned} \tag{2.8}$$

where the trace is over spinorial and configuration space indices.⁴ Integrating (2.8) by parts and expanding the logarithm in powers of the external potential finally gives

$$W = -\frac{1}{2\pi i} \int_C d\omega \, \text{Tr}_{\mathbf{x}} \log \left[\frac{\omega - H}{\omega - H_0} \right] \tag{2.9}$$

$$= \frac{1}{2\pi i} \sum_{k=1}^{\infty} \frac{1}{k} \int_C d\omega \, \text{Tr}_{\mathbf{x}} [\alpha G_0(\omega)]^k \tag{2.10}$$

Even with the free particle vacuum energy subtracted, Eq. (2.10) is still a formal, ultraviolet-divergent quantity. The culprit is the first term,⁵ $k = 2$, which is the familiar vacuum polarization diagram. This diagram is superficially quadratically divergent and has to be renormalized. However, because of gauge invariance, the actual ultraviolet divergence is milder. We will discuss this point in more detail in Section III. Let us simply mention here that

$$|W_{\text{ren}}^{(2)}| < R^3,$$

strictly. As R increases, the $k = 2$ term gives a vanishing contribution to the energy per unit volume of the star.⁶

The remaining contributions, $k \geq 4$, are ultraviolet convergent because the interaction term in (2.4) is renormalizable. On the other hand, because the neutrinos are massless, these terms have a strong infrared dependence

$$W^{(k)} \sim \frac{1}{k} \frac{1}{R} (\alpha R)^k. \tag{2.11}$$

For a neutron star, $\alpha \sim 20$ eV and $R \sim 10$ km, so that $\alpha R \sim 10^{12}$. With such an expansion parameter, the perturbative expansion of (2.1) is doomed to diverge and Eq. (2.10) must be resummed.

⁴The rule for closing the Feynman contour C in the complex ω plane in Eq. (2.8) is to take the average of the integral around the positive and negative energy cuts, so as to recover (2.2).

⁵Only the $k - \text{even}$ terms contribute for a static external potential. This is most easily seen in the Dirac representation of (2.4) because then the coupling to the potential involves the γ_5 matrix and the trace over an odd number of vertices is proportional to the Levi-Civita tensor. For a static potential, there are not enough indices to be contracted with the Levi-Civita tensor and the odd terms vanish identically.

⁶This statement is correct for a smooth effective potential, α . In a realistic star, the coarse grained structure at the neutron scale leads to a vacuum polarization contribution which scales like R^3 . See also the “Note added in proof.”

For comparison, the estimate of the vacuum energy (2.1) given in [5] is a large but finite sum,

$$W = \sum_{k=2}^N W^{(k)}, \quad (2.12)$$

where $N \sim 10^{57}$ is the number of neutrons in the star. Each term is taken to be given by the average k -body potential energy multiplied by the number of ways of choosing k neutrons from the grand total of N . For k large, but much less than N ,

$$W^{(k)} \sim \frac{G_F^k}{R^{2k+1}} \times \binom{N}{k} \sim \frac{1}{k!} \frac{1}{R} \left(\frac{G_F N}{R^2} \right)^k \sim \frac{1}{k!} \frac{1}{R} (\alpha R)^k. \quad (2.13)$$

Since $\alpha R \sim 10^{12}$, it is not surprising that after only a few terms the sum in Eq. (2.12) leads to an estimated energy which exceeds the mass of the neutron star. For future reference, let us quote the following estimate obtained in Ref. [8]. For fixed neutron density, corresponding to $\alpha \sim 20$ eV, a subsystem of radius $R \sim 2 \cdot 10^{-5}$ cm contains as much energy as the total mass of a neutron star with a radius of 10 km. Note also that $\alpha R = 20 \text{ eV} \times 2 \cdot 10^{-5} \text{ cm} \approx 20$.

Although similar, the estimates (2.11) and (2.13) differ crucially in the details. As emphasized by Abada, *et al* [9], for fixed k , there are $\mathcal{O}[(k-1)!]$ terms missing in (2.13). These are associated with multiple rescattering over the same neutrons. These processes also turn the finite sum (2.12) into an infinite series, as in (2.10). For large αR , the series is diverging and must be resummed.

B. The scattering problem

Consider the expression given in (2.2) for the vacuum energy. We can count the energy levels of the Hamiltonian (2.7) by putting the system in a large box [10]. Let us specify each eigenstate by its energy eigenvalue E and by a set of quantum numbers κ . For simplicity, consider a spherically symmetric potential in a large spherical box of radius R_{box} . (The parameter κ is then related to the total angular momentum.) Because the neutrinos are massless, the spectrum of (2.7) has no bound states but only scattering solutions,

$$\psi_{L\kappa} \sim \sin(kr + \phi_\kappa), \quad (2.14)$$

where k is the radial momentum. At the boundary of the box of radius R_{box} impose, for instance, that

$$k_{(0)} R_{\text{box}} + \phi_\kappa^{(0)} = n\pi \quad n = 0, 1, 2, \dots \quad (2.15)$$

where the index 0 refers to the free modes. From (2.15), the shift of energy level for fixed κ is then given by

$$E - E_0 \simeq \frac{dE}{dk} (k - k_0) = -\frac{1}{R_{\text{box}}} \frac{dE}{dk} (\phi_\kappa - \phi_\kappa^0) = -\frac{dE}{dk} \frac{\delta_\kappa}{R_{\text{box}}} \quad (2.16)$$

where δ_κ is the phase shift between the free and scattering eigenstates. Given that the number of modes per unit energy is

$$\frac{dn}{dE} = \frac{1}{\pi} R_{\text{box}} \frac{dk}{dE}, \quad (2.17)$$

we finally get

$$dn(E - E^0) = -\frac{1}{\pi} \delta_\kappa dE. \quad (2.18)$$

Hence, Eq. (2.2) becomes [10]

$$W = \frac{1}{2\pi} \sum_\kappa \int_0^\infty dE [\delta_\kappa(E) + \delta_\kappa(-E)]. \quad (2.19)$$

Of course, just as in (2.10), the expression Eq. (2.19) is a formal, ultraviolet-divergent quantity that must be renormalized.

Our strategy to address this issue is the following. (Details may be found in Section III, where we carry out an actual calculation.) Consider the expansion of the vacuum energy in the external potential of Eq. (2.10). As noted above, only the term of second order in the external potential is ultraviolet divergent. The evaluation of this term involves the vacuum polarization tensor $\Pi_{\mu\nu}$. Its divergent part can be absorbed using the standard renormalization procedure. The terms with $k \geq 4$ in the expansion (2.10) are ultraviolet-convergent. If $\alpha R \gg 1$ (with R the effective range of the potential), however, the terms with $k \geq 4$ in (2.10) must be resummed in order to get a sensible result. Using Eq. (2.10), this seems like a hopeless task.

Consider then the Taylor expansion of the phase shifts δ_κ in powers of the external potential αR . Only the first and second Born terms will lead to ultraviolet divergences in the expression of the vacuum energy (2.19). The first Born term is odd in E , and drops from the calculation if we integrate symmetrically over positive and negative energy phase shifts. If we subtract the second Born approximation from (2.19), the resulting, “renormalized vacuum energy” is free of ultraviolet divergences. The resulting quantity is equal to the resummed series of Eq. (2.10) with $k \geq 4$ and is precisely the quantity which has been estimated in Ref. [5].

In Section III we compute the renormalized vacuum energy for a spherical square well potential with depth α and radius R using Eq. (2.19). We numerically integrate the phase shifts over energy and subtract the second Born term. In principle we are required to evaluate all of the terms in (2.19). In practice, it is sufficient to investigate the convergence of the series in order to obtain a numerical estimate for the vacuum energy. As we will see, for large αR the number of terms to be calculated grows linearly with αR . Of course, we cannot expect to carry out such a calculation for an actual neutron star, for which $\alpha R \sim 10^{12}$. This is not necessary, however. Because the neutrinos are massless, there is just one dimensionless parameter, which is the effective expansion parameter αR . The question is then how the ground state energy changes from the domain where the perturbative expansion converges, $\alpha R \ll \mathcal{O}(1)$, to the domain where the expansion presumably breaks down because of the infrared divergence, $\alpha R \gg \mathcal{O}(1)$.

We have computed the renormalized vacuum energy for a number of points in the range $0 \leq \alpha R \leq 100$, including the point $\alpha R = 20$. According to Fischbach, *et al* [8], the energy corresponding to this point should be on the order of the mass of a neutron star,

$W \sim 10^{30} \text{ kg} \sim 10^{66} \text{ eV}$. By way of comparison, we find $W \simeq -2.3 \text{ keV}$. The energy is well-behaved over the whole range of our calculation. Remarkably, the energy per unit volume of the potential plotted as a function of αR exhibits a crossover at $\alpha R = \pi$. For $\alpha R > \pi$, the ground state of the system carries a non-zero neutrino charge.

The neutrino charge of the vacuum in the potential α is defined as

$$q = i \int d^3x \text{tr} [S_F(x, x') - S_F^0(x, x')]_{x' \rightarrow x} \quad (2.20)$$

$$\equiv -\frac{1}{2} \left\{ \sum_{i>0} - \sum_{i<0} \right\} \quad (2.21)$$

using the notation introduced in (2.2). Using Eqs. (2.15)-(2.18) gives (2.20) in terms of the phase shifts,

$$q = -\frac{1}{2\pi} \sum_{\kappa} \int_0^{\infty} dE \left\{ \frac{d\delta_{\kappa}(E)}{dE} - \frac{d\delta_{\kappa}(-E)}{dE} \right\}. \quad (2.22)$$

Just as was the case with the formal expressions for the ground state energy, the above expression for the vacuum charge will need to be renormalized. In this case this is accomplished by subtracting out the first Born term.

The existence of a neutrino condensate in the ground state is to be expected [7]. Because the neutrinos are massless, neutrino-antineutrino pairs are *a priori* easily produced by any non-zero gradient of the potential. Perhaps surprisingly, we have found that in our model it takes a finite value, $\alpha R = \pi$, for this to happen. For $\alpha R > \pi$, the ground state of our model contains a net neutrino charge. Note that the coupling term in (2.4) is precisely analogous to a constraint on the neutrino number with chemical potential $\mu = \alpha$ [18]. In the large volume limit, we expect the charge of the vacuum to be well approximated by that of a condensate with volume $V = 4\pi R^3/3$ [9],

$$q_{\text{cond}} = V \int \frac{d^3k}{(2\pi)^3} \theta(k - \alpha) = \frac{2(\alpha R)^3}{9\pi}. \quad (2.23)$$

Similarly, using the dispersion relation $\omega(k) = k - \alpha$ [which follows from (2.4) in the case of a homogeneous potential], the renormalized ground state energy should approach ⁷

$$W_{\text{cond}} = V \int \frac{d^3k}{(2\pi)^3} (k - \alpha) \theta(k - \alpha) = -\frac{\alpha^4 R^3}{18\pi} \quad (2.24)$$

That this is indeed the limiting behaviour of the renormalized vacuum energy for large potentials is demonstrated in Section III.

⁷In a neutron star, the induced neutrino charge density is extremely small, $\sim n_{\nu} \times 2 \cdot 10^{-23} \text{ fm}^{-3}$ (for $\alpha \sim 20 \text{ eV}$, and with n_{ν} the number of massless neutrino species), compared to the neutron density, $\sim 0.4 \text{ fm}^{-3}$. Consequently the back-reaction of the neutrino condensate on the neutron-induced effective potential is totally negligible. Also, because the low energy neutrinos of the condensate interact very weakly with each other, higher orders effects are presumably very small.

III. THE NEUTRINO GROUND STATE IN A SQUARE WELL POTENTIAL

A. Derivation of the phase shift

Both the charge and the energy of the neutrino ground state in the presence of a background potential may be expressed in terms of the scattering phase shifts. Our starting point, then, is the equation of motion following from the effective Lagrangian given in Eq. (2.4),

$$[\omega - i\vec{\sigma} \cdot \vec{\nabla} + \alpha(r)] \psi_L = 0, \quad (3.1)$$

where ψ_L is a two-component spinor, and where the potential is taken to be spherically symmetric. The energy, ω , may be positive or negative, with positive values corresponding to particle solutions and negative values corresponding to anti-particle solutions.

In order to solve for the scattering solutions we proceed, in the usual way [21,22], by decomposing them into angular momentum eigenstates,

$$\psi_{L_{jm}} = f_{jm}(r)\Omega_{jm}^{(+)} + g_{jm}(r)\Omega_{jm}^{(-)}, \quad (3.2)$$

where

$$\Omega_{jm}^{(+)} = (2l+1)^{-\frac{1}{2}} \begin{pmatrix} (l+m+\frac{1}{2})^{\frac{1}{2}} Y_l^{m-\frac{1}{2}} \\ (l-m+\frac{1}{2})^{\frac{1}{2}} Y_l^{m+\frac{1}{2}} \end{pmatrix}, \quad l = j - \frac{1}{2} \quad (3.3)$$

$$\Omega_{jm}^{(-)} = (2l'+1)^{-\frac{1}{2}} \begin{pmatrix} (l'-m+\frac{1}{2})^{\frac{1}{2}} Y_{l'}^{m-\frac{1}{2}} \\ - (l'+m+\frac{1}{2})^{\frac{1}{2}} Y_{l'}^{m+\frac{1}{2}} \end{pmatrix}, \quad l' = j + \frac{1}{2} > 0, \quad (3.4)$$

and where we use the standard notation Y_l^m for the spherical harmonics. The two-component spinors $\Omega_{jm}^{(\pm)}$ are eigenstates of J^2 and J_z , with eigenvalues $j(j+1)$ ($j=\frac{1}{2}, \frac{3}{2}, \dots$) and m , respectively. They are also eigenstates of L^2 , with orbital angular momentum $l=j-\frac{1}{2}$ for $\Omega_{jm}^{(+)}$ and $l'=j+\frac{1}{2}$ for $\Omega_{jm}^{(-)}$. In the following we will label the solutions by $l=0, 1, 2, \dots$ instead of by j , although it is to be understood that the solutions themselves are eigenstates of the total, not the orbital, angular momentum. With the above normalization for the angular momentum eigenstates we have the useful relation

$$\vec{\sigma} \cdot \hat{r} \Omega_{jm}^{(\pm)} = \Omega_{jm}^{(\mp)}. \quad (3.5)$$

Inserting (3.2) into (3.1), we then obtain a set of coupled first-order differential equations in f_l and g_l ,

$$\frac{df_l}{dr} - \frac{l}{r} f_l = -i[\omega + \alpha(r)] g_l \quad (3.6)$$

$$\frac{dg_l}{dr} + \frac{l+2}{r} g_l = -i[\omega + \alpha(r)] f_l. \quad (3.7)$$

As noted above, we will use a spherical square well potential in order to simplify our calculations. This choice has the benefit that the expression for the phase shift may be written out explicitly in terms of simple functions. We thus take

$$\alpha(r) = \alpha\theta(R - r), \quad r > 0. \quad (3.8)$$

The solutions of Eqs. (3.6) and (3.7) are then simply given by spherical Bessel functions. Requiring that the solutions be regular at the origin, and dropping an over-all normalization factor, we find

$$f_l = \begin{cases} j_l(|\omega + \alpha|r), & 0 < r < R \\ B [j_l(|\omega|r) \cos \delta_l + n_l(|\omega|r) \sin \delta_l], & r > R \end{cases} \quad (3.9)$$

$$g_l = \begin{cases} -i\varepsilon(\omega + \alpha)j_{l+1}(|\omega + \alpha|r), & 0 < r < R \\ -i\varepsilon(\omega)B [j_{l+1}(|\omega|r) \cos \delta_l + n_{l+1}(|\omega|r) \sin \delta_l], & r > R, \end{cases} \quad (3.10)$$

where B and δ_l are fixed by matching f_l and g_l at $r=R$ (note that the derivatives of f_l and g_l are in general discontinuous across the boundary) and where

$$\varepsilon(x) \equiv \frac{x}{|x|}. \quad (3.11)$$

Our conventions for the spherical Bessel functions are as follows:

$$j_l(\rho) = \left(\frac{\pi}{2\rho}\right)^{\frac{1}{2}} J_{l+\frac{1}{2}}(\rho), \quad n_l(\rho) = (-1)^l \left(\frac{\pi}{2\rho}\right)^{\frac{1}{2}} J_{-l-\frac{1}{2}}(\rho). \quad (3.12)$$

Performing the matching at the boundary, we finally arrive at the desired expression for the phase shift

$$\tan \delta_l(\omega) = \frac{\varepsilon(\omega)j_l(|\omega + \alpha|R)j_{l+1}(|\omega|R) - \varepsilon(\omega + \alpha)j_l(|\omega|R)j_{l+1}(|\omega + \alpha|R)}{\varepsilon(\omega + \alpha)n_l(|\omega|R)j_{l+1}(|\omega + \alpha|R) - \varepsilon(\omega)j_l(|\omega + \alpha|R)n_{l+1}(|\omega|R)}. \quad (3.13)$$

We may now use this expression in order to calculate the energy and charge of the vacuum.

B. Energy and charge of the vacuum

The formal expressions relating the energy and charge of the vacuum to the scattering phase shifts are given in (2.19) and (2.22). The spherical symmetry in our problem implies that the sum over κ becomes a sum over l . We may thus rewrite these expressions as

$$W = \frac{1}{2\pi} \sum_{l=0}^{\infty} (2l+2) \int_0^{\infty} d\omega [\delta_l(\omega) + \delta_l(-\omega)] \quad (3.14)$$

$$q = -\frac{1}{2\pi} \sum_{l=0}^{\infty} (2l+2) \int_0^{\infty} d\omega \left[\frac{\partial}{\partial \omega} \delta_l(\omega) - \frac{\partial}{\partial \omega} \delta_l(-\omega) \right]. \quad (3.15)$$

The factor $(2l+2) \equiv (2j+1)$ is the degeneracy factor for a given energy ω and total angular momentum j .

As we have discussed in Section II, the above formal expressions are in need of renormalization. Since our model is renormalizable, the divergences are confined to the first few

Born terms. The procedure to be followed in handling these divergences is then as follows: (i) Taylor-expand the formal expressions for the energy and the charge in order to isolate the divergences; (ii) subtract out the divergent terms and “resum” the Taylor expansion; (iii) renormalize the divergent terms in the usual way and add them back in. The resulting renormalized expressions are then finite and correspond to the actual energy and charge of the ground state of the system.

The first step in renormalizing the energy and the charge is to perform a Taylor expansion of the phase shift in αR . Let us then define

$$\delta_l(\omega; \alpha R) = (\alpha R)\delta_l^{(1)}(\omega) + (\alpha R)^2\delta_l^{(2)}(\omega) + (\alpha R)^3\delta_l^{(3)}(\omega) + \dots, \quad (3.16)$$

where

$$\delta_l^{(n)}(\omega) \equiv \frac{1}{n!} \left. \frac{\partial^n}{\partial z^n} \delta_l(\omega; z) \right|_{z \rightarrow 0}. \quad (3.17)$$

The explicit expressions for the first two Born terms are given by

$$\delta_l^{(1)}(\omega) = \varepsilon(\omega)\beta^2 \left\{ [j_l(\beta)]^2 + [j_{l+1}(\beta)]^2 - \frac{2(l+1)}{\beta} j_l(\beta) j_{l+1}(\beta) \right\}, \quad (3.18)$$

$$\begin{aligned} \delta_l^{(2)}(\omega) = & -\beta \left\{ \left[[j_l(\beta)]^2 + [j_{l+1}(\beta)]^2 - \frac{3(l+1)}{\beta} j_l(\beta) j_{l+1}(\beta) \right] \right. \\ & \left. - \beta^3 \left[n_l(\beta) j_l(\beta) + n_{l+1}(\beta) j_{l+1}(\beta) - \frac{1}{\beta} (l j_l(\beta) n_{l+1}(\beta) + (l+2) j_{l+1}(\beta) n_l(\beta)) \right] \right. \\ & \left. \times \left[[j_l(\beta)]^2 + [j_{l+1}(\beta)]^2 - \frac{2(l+1)}{\beta} j_l(\beta) j_{l+1}(\beta) \right] \right\}, \end{aligned} \quad (3.19)$$

where we have defined $\beta \equiv |\omega|R$. It is easy to convince oneself that the even (odd) Born terms are even (odd) in ω .

The formal expression for the energy is rendered finite by subtracting out the second Born term. (Since the first Born term is odd in ω , it drops out of the calculation if we integrate symmetrically over positive and negative energies.⁸) To compute the second Born term, we may use the expression for the $k = 2$ term in the expansion (2.10). Using the standard prescription, we may then regularize and renormalize this term. The renormalized expression for the energy is then given by

$$W_{\text{ren}} = W_{\text{ren}}^{(2)} + W^{(4+)}, \quad (3.20)$$

where

$$W^{(4+)} = \frac{1}{2\pi} \sum_{l=0}^{\infty} (2l+2) \int_0^{\infty} d\omega \left[\delta_l(\omega) + \delta_l(-\omega) - 2(\alpha R)^2 \delta_l^{(2)}(\omega) \right], \quad (3.21)$$

⁸Note that all terms in the Born expansion which are odd in α are also odd in ω and thus do not contribute to the energy. See the footnote following Eq. (2.10).

and where $W_{\text{ren}}^{(2)}$ denotes the renormalized vacuum polarization contribution. $W^{(4+)}$ corresponds to the resummed terms in the expansion (2.10) with $k \geq 4$.

The formal expression for the charge in (3.15) is divergent because the phase shifts tend to a constant as $\omega \rightarrow \infty$ [23],

$$\lim_{\omega \rightarrow \pm\infty} \delta_l(\omega) = \pm \int_0^\infty \alpha(r) dr = \pm \alpha R. \quad (3.22)$$

This divergence is removed by subtracting out the first Born term, which, as may easily be verified from (3.18), satisfies

$$\lim_{\omega \rightarrow \pm\infty} \delta_l^{(1)}(\omega) = \pm 1, \quad \delta_l^{(1)}(0) = 0. \quad (3.23)$$

Once regularized, this term gives a vanishing contribution to the renormalized charge (the charge is actually zero to all orders in perturbation theory), so that the final expression for the renormalized charge is given by

$$q_{\text{ren}} = \frac{1}{2\pi} \sum_{l=0}^{\infty} (2l+2) [\delta_l(0^+) - \delta_l(0^-)]. \quad (3.24)$$

The problem of calculating the charge of the vacuum then simply reduces to that of evaluating the particle and anti-particle phase shifts at the origin.

It is useful to note that for small αR the Taylor expansion of the phase shift given in (3.16) may be used to obtain a perturbative expansion for $W^{(4+)}$. For future reference, let us write this perturbative expansion as follows:

$$W^{(4+)} = \frac{1}{2\pi R} \sum_{n=4,6,8,\dots} (\alpha R)^n \sum_{l=0}^{\infty} (2l+2) I_l^{(n)}, \quad (\text{small } \alpha R) \quad (3.25)$$

where $I_l^{(n)}$ denotes the integral over ω of the n^{th} Born term,

$$I_l^{(n)} \equiv 2 \int_0^\infty d(\omega R) \delta_l^{(n)}(\omega). \quad (3.26)$$

This expansion is expected to be reliable for small αR . For large αR , of course, the perturbative expansion breaks down.

The main goal in the remainder of this paper will be to study the behaviour of the renormalized energy and charge as a function of αR . As we shall see, for fixed α both of these quantities scale like the volume of the potential region as the volume gets large. Before studying these quantities in detail, however, let us first consider the contribution to the total energy due to the vacuum polarization, $W_{\text{ren}}^{(2)}$. Since $W_{\text{ren}}^{(2)}$ scales roughly like the surface area of the potential region (for fixed α), its contribution to the total energy is eventually overwhelmed by that due to $W^{(4+)}$. Nevertheless, the handling of this term is somewhat subtle, so it is important to consider it carefully. Once we have studied $W_{\text{ren}}^{(2)}$, we will dispense with it altogether and consider only the contribution due to $W^{(4+)}$ when discussing the renormalized energy.

C. Vacuum polarization

We now turn to the calculation of $W_{\text{ren}}^{(2)}$. Consider the $k = 2$ term in the expansion (2.10),

$$W^{(2)} = \frac{1}{4\pi i} \int d\omega \int d^3x_1 d^3x_2 \text{tr} [\alpha(\vec{x}_1) G_0(\vec{x}_1, \vec{x}_2) \alpha(\vec{x}_2) G_0(\vec{x}_2, \vec{x}_1)]. \quad (3.27)$$

Using

$$G_0(\vec{x}) = - \int \frac{d^3k}{(2\pi)^3} \frac{e^{i\vec{k}\cdot\vec{x}}}{\omega - \vec{\sigma} \cdot \vec{k}}, \quad (3.28)$$

leads to

$$W^{(2)} = -\frac{1}{2i} \int d^3x_1 d^3x_2 \int \frac{d^3p}{(2\pi)^3} \alpha(\vec{x}_1) \alpha(\vec{x}_2) \Pi_{00}(\vec{p}) e^{i\vec{p}\cdot(\vec{x}_1 - \vec{x}_2)}, \quad (3.29)$$

where

$$\Pi_{\mu\nu}(p) = -\frac{i}{24\pi^2} \left(g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right) p^2 \left\{ \frac{2}{\epsilon} - \gamma + \log 4\pi + \frac{5}{3} - \log \left(\frac{-p^2}{\mu^2} \right) \right\} \quad (3.30)$$

is the familiar one-loop vacuum polarization tensor, which we have calculated using dimensional regularization. In Eq. (3.30), p is the 4-momentum in Minkowski space and μ is the renormalization scale. The divergent term, $\propto 1/\epsilon$, (plus an arbitrary constant term) can be absorbed as usual by redefining the parameters of the theory, $\alpha \rightarrow \alpha(\mu)$. Note that, because the polarization tensor is proportional to p^2 , the renormalized vacuum polarization term vanishes if the potential is homogeneous [9]. Introducing

$$\tilde{\alpha}(|\vec{p}|) = \alpha \int d^3x e^{i\vec{p}\cdot\vec{x}} \theta(R - r) = \frac{4\pi\alpha R^2}{|\vec{p}|} j_1(|\vec{p}|R), \quad (3.31)$$

with $j_1(x) = \sin(x)/x^2 - \cos(x)/x$, gives finally

$$W_{\text{ren}}^{(2)} = \frac{\alpha^2 R}{6\pi^2} \int_0^\infty dz z^2 j_1^2(z) \log \left(\frac{z^2}{(\mu R)^2} \right). \quad (3.32)$$

As it stands, the renormalized expression in (3.32) is still a linearly divergent quantity. We have chosen to work with a spherical square well potential and, because of the presence of the sharp edge, arbitrarily high radial momenta contribute to the integral. Any smoother potential would lead to a finite result. At the level of the effective theory, there is a natural ultimate cut-off provided by the size of the neutron, $\Lambda \sim 0.2$ GeV. (This also suggests the choice $\mu = \Lambda$.) Inserting the cut-off, we get that the vacuum polarization term (3.32) scales roughly like the area of the potential, $W_{\text{ren}}^{(2)} \sim \alpha^2 R^2 \Lambda$. Consequently, $W_{\text{ren}}^{(2)}/R^3$ contributes a vanishing amount to the energy density as the size of the system increases. Let us emphasize that the remainder of our calculation is completely insensitive to the presence of this cut-off since the higher order terms in (2.10) are ultra-violet convergent.

D. Numerical evaluation of the renormalized energy and charge

We turn now to a numerical evaluation of the renormalized energy and charge. From now on we will ignore the vacuum polarization contribution to the energy. We will thus loosely refer to $W^{(4+)}$ as the renormalized energy.

1. Small values of αR

Let us begin our study of the charge and energy of the ground state by considering rather modest values of αR . Fig. 1(a) shows plots of the particle and antiparticle phase shifts (solid curves) as functions of ω for $l = 0$ in the case that $\alpha R = 1$. Also shown are the sums of the first and second Born approximations to the phase shifts (dashed curves). Note that the particle (antiparticle) phase shifts approach $+(-)\alpha R$ as $\omega \rightarrow \infty$. Furthermore, the phase shifts go to zero at the origin so that, according to Eq. (3.24), the vacuum charge is zero.⁹ The plots for higher values of l are qualitatively similar.

Fig. 1(b) shows a plot – again for $\alpha R = 1$ and $l = 0$ – of the sum of the particle and antiparticle phase shifts with the second Born term subtracted. This is the quantity which must be integrated (and summed over l) in order to obtain the renormalized energy in Eq. (3.21).

For small enough values of αR , one expects the energy to be well-described by the leading term in the perturbative expansion defined in (3.25). Our numerical results show that this is indeed the case for $\alpha R \lesssim 1/2$, which actually provides a non-trivial check on our results. The advantage of using the perturbative expansion to evaluate the renormalized energy for small αR is that the integrals over ω may be done exactly in that case,¹⁰ whereas those for the exact expression, (3.21), must be done numerically. Furthermore, in the case of the exact expression, the numerical integrations over ω become increasingly difficult as $\alpha R \rightarrow 0$.

The leading term in the perturbative expansion is at fourth order in αR . Fig. 2 shows a log-log plot of $(-1) \times (2l + 2) \times I_l^{(4)}$ as a function of l . (We have not plotted the point

⁹ It may at first seem surprising that the charge vanishes for $\alpha R = 1$. After all, the neutrinos are massless and it would seem that any gradient of the potential could produce neutrino-antineutrino pairs. However, this argument neglects the effect of the neutrino's spin. To see this, consider the system of first-order differential equations for f_l and g_l , (3.6) and (3.7). Writing these as two decoupled second-order Schrödinger equations shows that the g_l component always has a repulsive centrifugal barrier,

$$\begin{aligned} \frac{d^2 f_l}{dr^2} + \frac{2}{r} \frac{df_l}{dr} - \frac{l(l+1)}{r^2} f_l + (\omega + \alpha)^2 f_l &= 0 \\ \frac{d^2 g_l}{dr^2} + \frac{2}{r} \frac{dg_l}{dr} - \frac{(l+1)(l+2)}{r^2} g_l + (\omega + \alpha)^2 g_l &= 0 \end{aligned}$$

Thus, even for the $l = 0$ mode, it takes a finite αR to create a charge in the vacuum.

¹⁰We do not include the explicit form of $\delta_l^{(4)}(\omega)$ here because it is rather unwieldy. As was the case for $\delta_l^{(2)}(\omega)$, however, the expression for $\delta_l^{(4)}(\omega)$ may be expressed in terms of spherical Bessel functions. Thus the integrals may be done analytically (albeit with the aid of a computer).

corresponding to $l = 0$ for obvious reasons.) The sum over l of this quantity is directly proportional to the renormalized energy in the perturbative region. As is clear from the figure, this quantity has roughly a power-law fall-off as $l \rightarrow \infty$. Using the slope of the curve in Fig. 2, we find that $(2l + 2) \times I_l^{(4)} \propto l^{-1.9}$, for large l . This allows us to obtain an approximate answer for the sum over l ,

$$\sum_{l=0}^{\infty} (2l + 2) I_l^{(4)} = -0.00721 \pm 0.00005, \quad (3.33)$$

where the quoted uncertainty is a rather conservative estimate of the error incurred in the extrapolation to large l . We are thus able to compute the leading perturbative contribution to $W^{(4+)}$, which is given by

$$W^{(4+)} = -0.00115(\alpha R)^4/R + \dots \quad (3.34)$$

2. Larger values of αR

As αR is increased, one finds that there is a critical point beyond which it becomes energetically favourable for the neutrinos to condense; thus, for $\alpha R > \alpha R|_{\text{crit}}$, the ground state of the system contains a neutrino condensate. The presence of the condensate has a very interesting effect on the renormalized energy. Before calculating the energy, however, let us first examine the phase shifts for large αR . Fig. 3 shows plots of the particle and antiparticle phase shift for $\alpha R = 10$, with $l = 0, 2, 4$ and 6 . These plots are qualitatively very different from those for small αR [c.f. Fig. 1(a)] in that the phase shift in the strong-potential case exhibits resonances. Furthermore, for l less than some critical value the phase shifts do not vanish at the origin, as they do in the weak-potential case.¹¹ Thus, according to Eq. (3.24), the ground state is charged:

$$q_{\text{ren}}(\alpha R = 10) = \sum_{l=0}^{l_{\text{max}}} q_l = 56. \quad (3.35)$$

Note that the sum over l for the charge always truncates at some l_{max} . For $\alpha R = 10$, $l_{\text{max}} = 5$.

The charge of the ground state as a function of αR is intimately tied to the resonance structure of the particle and antiparticle scattering solutions. It is well-known in non-relativistic scattering theory that as a potential well is deepened, the resonances of the scattering solutions migrate to the origin of the complex k -plane and, at some critical strength of the potential, join the imaginary axis and become true bound states. As the potential is made deeper and deeper, the bound states move to lower and lower energies (i.e., they become more and more tightly bound). The situation is slightly different in the relativistic

¹¹Note, however, that Levinson's theorem is still satisfied: since there are no bound states, the sum of the particle and anti-particle phase shifts at the origin is zero [23].

scattering theory for fermions. Consider first the case in which the fermion has a non-zero mass. In that case, the resonances of the positive energy (particle) scattering solutions can migrate to the imaginary axis in the k -plane and become true bound states, just as in the non-relativistic case. As the potential is deepened still more, however, the bound states eventually turn around and re-enter the continuum, this time as resonances for the negative energy (antiparticle) scattering solutions [14]. The presence of resonances in the negative energy scattering solutions signals that the ground state has become charged. The situation is similar in the massless case, except that in this case the positive energy resonances move to the origin and pass immediately to the negative energy continuum without ever making true bound states.

In our simple model, it is possible to calculate precisely the critical values of the potential at which the charge of the ground state changes discontinuously. The solutions of the “zero-energy resonances” of the system are normalizable solutions of the Weyl equation.¹² Setting ω to zero in Eqs. (3.6) and (3.7), we find that f_l and g_l decouple for $r > R$, giving

$$f_l \propto r^l \quad (3.36)$$

$$g_l \propto r^{-(l+2)}, \quad (3.37)$$

for $r > R$. The normalizable solutions, then, are those which kill f_l in the asymptotic region. After a few lines of algebra, one arrives at the following simple condition for the critical parameters at which the charge of the vacuum changes:

$$j_l(\alpha R) = 0. \quad (3.38)$$

In particular, then, the vacuum goes from being uncharged to having charge equal to 2 when $\alpha R = \pi$.

Using the relation (3.38), it is straightforward to calculate the charge of the ground state for any value of αR : for each l , one simply needs to count the number of zeros of $j_l(\rho)$ with $\rho < \alpha R$. Fig. 4 shows a plot of the charge of the ground state as a function of αR . It is clear from the solid curve in this plot that the charge does indeed change discontinuously as αR is increased. The dashed curve in this plot shows the charge which one obtains in the large-volume limit for a system with chemical potential $\mu = \alpha$, Eq. (2.23). As αR is increased, we find that our exact result tends to this value.

Let us now calculate the renormalized energy, $W^{(4+)}$, for $\alpha R > \pi$. Let us define

$$W^{(4+)} \equiv \sum_{l=0}^{\infty} W_l^{(4+)}, \quad (3.39)$$

where

$$W_l^{(4+)} = \frac{1}{2\pi}(2l+2) \int_0^{\infty} d\omega \left[\delta_l(\omega) + \delta_l(-\omega) - 2(\alpha R)^2 \delta_l^{(2)}(\omega) \right] \quad (3.40)$$

$$= \frac{1}{2\pi}(2l+2) \int_0^{\infty} d\omega \left[\delta_l(\omega) + \delta_l(-\omega) \right] - \frac{(\alpha R)^2}{2\pi R}(2l+2) I_l^{(2)} \quad (3.41)$$

¹²These are the “would-be” bound states of the system which, in a vector-like theory with a massive fermion, would have migrated onto the imaginary axis in the k -plane and become bound states.

In the second line we have explicitly separated out the term corresponding to the integral of the second Born term, $I_l^{(2)}$, since it may be calculated analytically. (See Fig. 5.) We stress that for a given l both terms in (3.41) are finite.

Fig. 6 shows two plots of the dimensionless combination $W_l^{(4+)}R$ as a function of l for two strong potentials. In each plot the solid dots correspond to $\alpha R = 10$ and the open circles correspond to $\alpha R = 20$. These plots differ qualitatively from the analogous plots in the case of weak potentials. In the present case, the energy contains a large “bump” for $l \leq l_{\text{crit}}$ (see Fig. 2 for comparison). For $l > l_{\text{crit}}$, the energy has a power-law fall-off which is similar to the large- l behaviour found in the weak-potential case. From the slope of the tail in the log-log plot, one sees that for large l the tail goes roughly as l^{-2} .¹³ The presence of the bump in the plot of $W_l^{(4+)}$ is directly related to the presence of a condensate. In all cases which we have studied, the value of l at which the energy suddenly drops is precisely the same value of l for which there is no more contribution to the vacuum charge; *i.e.* $l_{\text{crit}} = l_{\text{max}} \sim \alpha R$.

The final step in our calculation is to perform the sum of $W_l^{(4+)}$ over l in order to obtain the renormalized energy, and then to study the behaviour of the renormalized energy as a function of αR . Fig. 7 contains plots of both the renormalized charge and the renormalized energy as functions of the dimensionless parameter αR . This figure represents the central result of our paper. The plots of both the charge and the energy have been normalized to the values which one obtains assuming a condensate with chemical potential $\mu = \alpha$, Eq. (2.23) and Eq. (2.24).

For $\alpha R < \pi$, the charge is equal to zero and the renormalized energy is well described by keeping only the first non-vanishing term in the perturbative expansion [see Eq. (3.34)]. This limit is indicated by the dashed line in Fig. 7(b). For $\alpha R > \pi$ the energy is no longer well-described by the leading term in the perturbative expansion, and instead approaches the value which one expects for a large-volume system with chemical potential $\mu = \alpha$.

Fig. 7 demonstrates very convincingly that the magnitude of the shift in energy of the vacuum does not increase exponentially as the potential becomes large, but rather crosses over smoothly to the value expected for a condensate of massless neutrinos. The numerical calculations become quite computer-intensive for large αR and we have not gone beyond $\alpha R = 100$. In our opinion, there is no compelling reason to go to larger values of αR .

IV. CONCLUSIONS

In this paper we have presented an exact non-perturbative one-loop calculation of the neutrino vacuum energy in the presence of an external electroweak potential. In order to simplify the calculation, we have chosen to work with a spherical square well potential with depth α and radius R .

¹³Note that a very precise determination (to better than one part in 10^5) of the two terms contributing to $W_l^{(4+)}$ in Eq. (3.41) is required in order to see that the tail in Fig. 6(b) is actually converging.

The formal expression for the vacuum energy is ultraviolet divergent. This divergence is however limited to the leading term in the expansion of the vacuum energy in powers of αR and can be renormalized using the standard methodology. The higher order terms are free of ultraviolet divergences. For large values of αR , however, these terms are infrared divergent. The non-perturbative resummation of these terms has been the main goal of our paper. We refer to the resummed expression as the renormalized vacuum energy.

The perturbative expansion of the renormalized vacuum energy is reliable for small values of αR . For $\alpha R > \pi$, the perturbative expansion breaks down and we find that the ground state contains a non-zero neutrino charge. The neutrino charge may be calculated exactly for any αR in our model. As αR increases, the charge is well-approximated by the charge of a neutrino condensate with chemical potential α . The onset of the charged vacuum at $\alpha R = \pi$ is accompanied by a smooth cross-over in the vacuum energy. (See Fig. 7.) For large αR , the energy approaches the value expected for a neutrino condensate.

It has been argued in Refs. [5,8] that long range forces due to the exchange of massless neutrinos could destabilize a neutron star. Given that a finite density of neutrons gives rise to an effective electroweak potential for the low energy neutrinos we can directly compare our results with these claims. According to [8], a dense microscopic system of neutrons with radius $R \sim 2 \cdot 10^{-5}$ cm and $\alpha \sim 20$ eV contains as much energy due to the exchange of massless neutrinos as the total mass of a neutron star of radius 10 km; *i.e.*, $W \sim 10^{66}$ eV. By way of contrast, the infrared-sensitive terms in our calculation contribute $W \sim -2.3$ keV per massless neutrino species. As emphasized previously by Abada, *et al* [9], the erroneous results obtained in [5,8] are due to an improper counting of the processes contributing to the vacuum energy. When properly resummed, these processes lead to small effects, even for massless neutrinos, and do not endanger a neutron star.

NOTE ADDED IN PROOF

In this note we address some issues raised in two manuscripts [24,25] which appeared after the completion of the present work.

In Ref. [24], Fischbach and Woodahl have questioned the validity of the effective field theory approach used in the present paper. Their criticism is based on a discrepancy between our estimate of the two-body contribution to the vacuum energy of a neutron star [see the discussion below Eq. (3.32)],

$$W^{(2)} \sim \alpha^2 R^2 \Lambda, \quad (4.1)$$

and their estimate [Eqs. (22) and (23) in [24]],

$$W^{(2)} \sim \alpha^2 R^3 \Lambda^2. \quad (4.2)$$

It is however straightforward to reconcile these estimates. The discrepancy may be traced to the fact that we have used a smooth potential to model the star. This mean field approximation, while very well-suited for studying the infrared behavior of the system, is a poor approximation to a realistic star in the ultraviolet, where the coarse grained structure becomes important. It is easy to verify that upon introducing fluctuations on scales of order

$1/\Lambda$ and repeating the analysis of Section III.C, one obtains an estimate for $W^{(2)}$ which is in qualitative agreement with Eq. (4.2).

We reiterate, however, that the higher-point contributions to the vacuum energy [i.e., the terms with $k \geq 4$ in Eq. (2.10)] are insensitive to the ultraviolet features of the star. As discussed at length in the text, these contributions must be resummed, a task for which the effective field theory approach in the mean field approximation is correct and perfectly suited. The resummed result, $W^{(4+)}$, tends to the value expected for a condensate of neutrinos with chemical potential α as αR gets large. This quantity gives the leading contribution to the vacuum energy for a smooth potential. For a realistic star, the dominant contribution to the vacuum energy is likely to be due to the two-point contribution.

In Ref. [25], Abada, Pène and Rodriguez-Quintero have independently analyzed the behavior of massless neutrinos in a three-dimensional electroweak potential and have, like us, concluded that the exchange of massless neutrinos does not threaten the stability of a neutron star. However, their estimate of the vacuum energy,

$$W \sim \alpha R^3 \Lambda^3, \quad (4.3)$$

differs both from our two-body contribution (4.1), valid for a smooth potential, and from our estimate of the contribution due to the higher-point diagrams,

$$W^{(4+)} \simeq W_{\text{cond}} = -\frac{\alpha^4 R^3}{18\pi}. \quad (4.4)$$

We believe that the estimate in Eq. (4.3) is incomplete. Note that this expression is linear in the external potential α . Changing the sign of the external potential is, however, equivalent to considering an antineutron star (which attracts antineutrinos) instead of a neutron star (which attracts neutrinos). This transformation leaves the vacuum energy unchanged. That the vacuum energy is an even function of α is indeed manifest to all orders in the Schwinger-Dyson expansion. In particular, the tadpole diagram [which behaves superficially like the expression in Eq. (4.3)] vanishes identically when both the positive and negative energy eigenstates are taken into account in the summation over modes. The first non-vanishing contribution is then the quadratic term as discussed above.

ACKNOWLEDGMENTS

We wish to thank M. Gyulassy, S. Nussinov, F. Paige, R.D. Pisarski, J. Sapirstein and N. Weiss for useful comments. We are particularly indebted to R. Jaffe for many insightful and helpful conversations. This research was supported in part by the U.S. Department of Energy under contract number DE-AC02-76CH00016. K.K. is also supported in part by the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

- [1] G. Feinberg and J. Sucher, Phys. Rev. **166**, 1638 (1968); G. Feinberg and J. Sucher and C.-K. Au, Phys. Rept. **180**, 83 (1989).
- [2] S.D.H. Hsu, P. Sikivie, Phys.Rev. **D49**, 4951 (1994); J.A. Grifols, E. Masso and R. Toldra, Phys.Lett. **B389**, 563 (1996).
- [3] R.P. Feynman, F.B. Morinigo and W.G. Wagner, Feynman lectures on Gravitation, Addison-Wesley, Readings, MA (1995).
- [4] J.B. Hartle, Phys. Rev. **D1**, 394 (1970).
- [5] E. Fischbach, Ann. of Phys. **247**, 213 (1996).
- [6] A. Y. Smirnov and F. Vissani, hep-ph/9604443.
- [7] A. Loeb, Phys. Rev. Lett. **64** (1990) 115.
- [8] B. Woodahl, M. Parry, S.-J. Tu, E. Fischbach, hep-ph/9709334.
- [9] As. Abada, M.B. Gavela and O. Pène, Phys. Lett. B **387**, 315 (1996).
- [10] J. Schwinger, Phys. Rev. **94**, 1362 (1954).
- [11] R. Dashen, B. Hasslacher, A. Neveu, Phys. Rev. **D10**, 4114 (1974); *ibid*, 4130 (1974).
- [12] E. Farhi, N. Graham, P. Haagsen and R.L. Jaffe, “Finite quantum fluctuations about static field configurations,” hep-th/9802015.
- [13] M. Bordag and K. Kirsten, Phys. Rev. **D53**, 5753 (1996).
- [14] W. Greiner, B. Müller and J. Rafelski, *Quantum electrodynamics of strong fields*, Berlin, Springer (1985).
- [15] As. Abada, O. Pène and J. Rodriguez-Quintero, “Multibody neutrino exchange in a neutron star: neutrino sea and border effects,” hep-ph/9712266.
- [16] M. Kachelriess, “Neutrino self-energy and pair creation in neutron stars,” hep-ph/9712363.
- [17] P.D. Mannheim, Phys. Rev. **D 37**, 1935 (1988); P.B. Pal and T.N. Pham, Phys. Rev. **D 40** (1989) 259; L.N. Chang and R.K.P. Zia, Phys. Rev. **D 38** (1988) 1669.
- [18] K.A. Kiers, Ph.D. thesis, University of British Columbia, 1996.
- [19] K. Kiers, N. Weiss, Phys. Rev. **D56**, 5776 (1997), hep-ph/9704346.
- [20] L. Wolfenstein, Phys. Rev. **D 17** (1978) 2369; Phys. Rev. **D 20** (1979) 2634; S.P. Mikheyev and A.Yu. Smirnov, Yad. Fiz. **42** (1985) 1441 [Sov. J. Nucl. Phys. **42** (1985) 913]; Il Nuovo Cimento **C 9** (1986) 17.
- [21] C. Itzykson and J.-B. Zuber, *Quantum Field Theory*, McGraw-Hill, NY (1980).
- [22] A. Messiah, *Quantum Mechanics*, Vol. II, John Wiley & Sons, NY.
- [23] Z.-Q. Ma and G.-J. Ni, Phys. Rev. **D D31**, 1482 (1985).
- [24] E. Fischbach and B. Woodahl, “Neutrino exchange interactions in one-dimension, two-dimensions, and three-dimensions,” hep-ph/9801387.
- [25] As. Abada, O. Pène and J. Rodriguez-Quintero, “Finite size effects on multibody neutrino exchange,” hep-ph/9802393.

FIGURES




FIG. 1. Plots of the phase shift for $\alpha R = 1$ and $l = 0$. (a) The upper (lower) solid curves give the particle (antiparticle) phase shifts, while the dashed curves give the sums of the first and second Born terms. (b) This curve shows the sum of the particle and antiparticle phase shifts with twice the second Born term subtracted.




FIG. 2. Plot of $-(2l+2)I_l^{(4)}$ vs. l .




FIG. 3. Plots of the particle and antiparticle phase shifts for $\alpha R = 10$ and $l = 0, 2, 4$ and 6 . In each case the upper curve is the particle phase shift and the lower curve is the antiparticle phase shift.




FIG. 4. Plot of the charge as a function of αR . The solid curve gives the exact charge, which has periodic jumps, and the dashed curve gives the charge expected for a condensate in the large-volume limit.




FIG. 5. Plot of $I_l^{(2)}$ vs. l .




FIG. 6. Plots of $-W_l^{(4+)}R$ vs. l for $\alpha R = 10$ (solid dots) and $\alpha R = 20$ (open circles). The slopes of the tails in the log-log plots are approximately -2 in each case.




FIG. 7. Plots of (a) the charge and (b) the energy as a function of αR . The dots give the results of our exact (non-perturbative) calculations. Both the charge and the energy are normalized to the contribution expected from a condensate in the large-volume limit. The lower dashed curve in (b) gives the value obtained in the limit of small αR (see Eq. (3.34)).